FRAMINGHAM RISK SCORE (FRS)

Estimation of 10-year Cardiovascular Disease (CVD) Risk

Date: _____

Patient's Name: _____

Step 1[1]

In the "points" column enter the appropriate value according to the patient's age, HDL-C, total cholesterol, systolic blood pressure, and if they smoke or have diabetes. Calculate the total points.

Risk Factor		Risk Points		Points
		Men	Women	
Age				
30-34		0	0	
35-39		2	2	
40-44		5	4	
45-49		7	5	
50-54		8	7	
55-59		10	8	
60-64		11	9	
65-69		12	10	
70-74		14	11	
75+		15	12	
HDL-C (mmol/L)				
>1.6		-2	-2	
1.3-1.6		-1	-1	
1.2-1.29		0	0	
0.9-1.19		1	1	
<0.9		2	2	
Total Cholesterol				
<4.1		0	0	
4.1-5.19		1	1	
5.2-6.19		2	3	
6.2-7.2		3	4	
>7.2		4	5	
Systolic Blood Pressure (mmHg)		Not Treated / Treated	Not Treated / Treated	
<120		-2 / 0	-3 / -1	
120-129		0 / 2	0 / 2	
130-139		1 / 3	1 / 3	
140-149		2 / 4	2 / 5	
150-159		2 / 4	4 / 6	
160+		3 / 5	5 / 7	
Smoker	Yes	4	3	
	No	0	0	
Diabetes	Yes	statin-indicated condition		
	No	0	0	
Total Points				

1 Adapted from: D'Agostino RB et al.(i). General cardiovascular risk profile for use in primary care. The Framingham Heart Study. Circ 2008;117:743-53.
2 Adapted from: Genest J et al.(i). 2009 Canadian Cardiovascular Society/Canadian guidelines for the diagnosis and treatment of dyslipidemia and prevention of cardiovascular disease in the adult. Can J Cardiol. 2009;25(10):567-579.
3 Adapted from: Anderson T et al.(i). 2012 Update of the Canadian Cardiovascular Society guidelines for the diagnosis and treatment of dyslipidemia for the prevention of cardiovascular disease in the adult. Can J Cardiol. 2013;29(2):151-167.

‡ apoB: apolipoprotein B stat, CVD: cardiovascular disease, FRS: Framingham Risk Score, HDL-C: high-density lipoprotein cholesterol, LDL-C: low-density lipoprotein cholesterol.
* Statins indicated as initial therapy
** Consider LDL-C < 1.8 mmol/L for subjects with acute coronary syndrome (ACS) within past 3 months

Provided courtesy of **Canadian Cardiovascular Society**
Leadership. Knowledge. Community.

Step 2[1]

Using the total points from Step 1, determine the 10-year CVD risk (%).*

Total Points	10-Year CVD Risk (%)*	
	Men	Women
-3 or less	<1	<1
-2	1.1	<1
-1	1.4	1.0
0	1.6	1.2
1	1.9	1.5
2	2.3	1.7
3	2.8	2.0
4	3.3	2.4
5	3.9	2.8
6	4.7	3.3
7	5.6	3.9
8	6.7	4.5
9	7.9	5.3
10	9.4	6.3
11	11.2	7.3
12	13.3	8.6
13	15.6	10.0
14	18.4	11.7
15	21.6	13.7
16	25.3	15.9
17	29.4	18.51
18	>30	21.5
19	>30	24.8
20	>30	27.5
21+	>30	>30

Step 3[1]

Using the total points from Step 1, determine heart age (in years).

Heart Age, *y*	Men	Women
<30	<0	<1
30	0	
31		1
32	1	
34	2	2
36	3	3
38	4	
39		4
40	5	
42	6	5
45	7	6
48	8	7
51	9	8
54	10	
55		9
57	11	
59		10
60	12	
64	13	11
68	14	12
72	15	
73		13
76	16	
79		14
>80	≥17	15+

* Double cardiovascular disease risk percentage for individuals between the ages of 30 and 59 without diabetes if the presence of a positive history of premature cardiovascular disease is present in a first-degree relative before 55 years of age for men and before 65 years of age for women. This is known as the modified Framingham Risk Score.[3]

Step 4[2,3]

Using 10-year CVD risk from Step 2, determine if patient is Low, Moderate or High risk.† Indicate Lipid and/or Apo B targets

Risk Level†	Initiate Treatment If:	Primary Target (LDL-C)	Alternate Target
High FRS ≥20%	• Consider treatment in all *(Strong, High)*	• ≤2 mmol/L or ≥50% decrease in LDL-C *(Strong, Moderate)*	• Apo B ≤0.8 g/L or • Non-HDL-C ≤2.6 mmol/L *(Strong, High)*
Intermediate FRS 10-19%	• LDL-C ≥3.5 mmol/L *(Strong, Moderate)* • For LDL-C <3.5 mmol/L consider if: • Apo B ≥1.2 g/L • OR Non-HDL-C ≥4.3 mmol/L *(Strong, Moderate)* • *Men ≥50 and women ≥60 with 1 risk factor: low HDL-C, impaired fasting glucose, high waist circumference, smoker, hypertension*	• ≤2 mmol/L or ≥50% decrease in LDL-C *(Strong, Moderate)*	• Apo B ≤0.8 g/L or • Non-HDL-C ≤2.6 mmol/L *(Strong, Moderate)*
Low FRS <10%	• statins generally not indicated	• statins generally not indicated	• statins generally not indicated
Statin-indicated conditions**	• *Clinical atherosclerosis** • *Abdominal aortic aneurysm* • *Diabetes mellitus Age ≥ 40 years 15-Year duration for age ≥ 30 years (DM1) Microvascular disease* • *Chronic kidney disease (age ≥ 50 years) eGFR <60 mL/min/1.73 m2 or ACR > 3 mg/mmol*		

Lipid targets LDL-C: _____ or Apo B: _____